

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C11B 3/14, A23D 7/00, 9/00	A1	(11) International Publication Number: WO 99/32589 (43) International Publication Date: 1 July 1999 (01.07.99)
(21) International Application Number: PCT/EP98/07797 (22) International Filing Date: 1 December 1998 (01.12.98) (30) Priority Data: 97204048.9 19 December 1997 (19.12.97) EP 98200126.5 19 January 1998 (19.01.98) EP (71) Applicant (for all designated States except AU BB CA CY GB GD GH IE KE LK LS MN MW NZ SD SG SZ TT UG): UNILEVER N.V. [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL). (71) Applicant (for AU BB CA CY GB GD GH IE KE LK LS MN MW NZ SD SG SZ TT UG only): UNILEVER PLC [GB/GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB). (72) Inventors: VAN BUUREN, Jan; Unilever Research Vlaardingen, Olivier van Noortlaan 120, NL-3133 AT Vlaardingen (NL). GANGULI, Keshab, Lal; Unilever Research Vlaardingen, Olivier van Noortlaan 120, NL-3133 AT Vlaardingen (NL). VAN PUTTE, Karel, P., A., M.; Unilever Research Vlaardingen, Olivier van Noortlaan 120, NL-3133 AT Vlaardingen (NL).		(74) Agent: SIKKEN, Antonius, H., J., M.; Unilever N.V., Patent Dept., P.O. Box 137, NL-3130 AC Vlaardingen (NL). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: OLIVE OIL CONTAINING FOOD COMPOSITION (57) Abstract A method has been found for the manufacture of a spread which is characterized by the presence of a substantial amount of olive oil polyphenols and the absence of olive oil odour. The deodorized olive oil to be used for the preparation is obtained by a mild refining process which spares the olive oil's minor components.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

OLIVE OIL CONTAINING FOOD COMPOSITION

5 The present invention deals with a fat containing food composition, particularly a spread, which contains a substantial amount of olive oil. Such composition is believed to fit in a healthy diet particularly by helping to control blood cholesterol content. The invention also
10 comprises a process for preparation of the composition.

BACKGROUND OF THE INVENTION

15 The present invention deals with a food composition with a substantial fat content. Since fat consumption is associated with an increased risk of cardiovascular disorders, the nutritional value of different types of fat as well as methods for reducing the fat content of fat
20 compositions had been the object of extensive investigations. Presently not only the triglyceride part of fats is the subject of investigations, but also the nature and the effects on health of fat attributes, the so-called minor components which are present in small amounts in non-
25 refined natural fats. It has recently been found that the minor components which are denoted as anti-oxidants, including fat polyphenols, positively interfere with the body's cardiovascular system.

Traditionally, natural fats are refined before they are
30 used as an ingredient for the preparation of food. However, traditional fat refining is not discriminating to the nature of the fat ingredients and aims at the removal of

all substances other than triglycerides, including minor components.

Spreads are food compositions which usually contain a substantial amount of fat, often 40 wt.% or more. Usually the fat consists of a liquid oil and a structuring fat which gives the fat blend a proper consistency. Olive oil is a valuable nutritional fat because of its high content of oleic acid. Olive oil has been incorporated in food compositions either unrefined or fully refined.

In EP 421 504 spreads are described in which the fat phase consists partially of unrefined olive oil. According to GB 2281304 refined olive oil is used in the liquid oil blend of the fat phase for spread manufacture. Even when used in small amounts, the incorporation of unrefined olive oil imparts a much pronounced flavour and taste. The characteristic olive oil flavour often is appreciated, but by many people this flavour occurring in spreads is perceived as offensive and therefore is unsuitable for spreads. Hence the olive oil has to be deodorised by a refining treatment. Olive oil traditionally is deodorized at temperatures of 250°C or even more. Under such drastic conditions also the oil's minor components, including natural anti-oxidants, particularly the typical olive oil polyphenols such as oleuropein, its aglycon (shortly aglycon), tyrosol and hydroxytyrosol, are nearly fully stripped away.

WO 97/06697 teaches that food products can be fortified with anti-oxidants, including polyphenols and particularly the water soluble tea polyphenols.

Polyphenols present in olive oil are more or less fat soluble and have a chemical composition which is different from the water soluble polyphenols. They are presumed to play in the body an anti-oxidant role in specific tissues, 5 possibly because of their fat solubility.

When incorporating a substantial amount of olive oil polyphenols in spreads, imparting a strong olive oil odour could not be avoided up to now.

10

SUMMARY OF THE INVENTION

We have found an olive oil containing spread which contains 15 a considerable amount of olive oil polyphenols but which nevertheless is free from any offensive olive oil odour. The invention makes use of a process for mild olive oil refining which allows largely retaining the polyphenols. With this process the prejudice has been overcome that 20 proper olive oil refining requires drastic refining conditions with regard to temperature and process time. Traditional refining removes free fatty acids, resulting in a content of less than 0.1 wt.%. The found refining conditions relating to flow of sparging gas, temperature, 25 pressure and process time are less drastic, but nevertheless are appropriate for the removal of olive oil odour and - to a large extent - of fatty acids, so that the refined olive oil is pure enough to be used for the manufacture of a spread, while the process spares a major 30 part of the minor components, including the olive oil polyphenols, tocopherols, squalene and further unsaponifiables, including fatty alcohols and sterols, many

of which are believed to be valuable nutrients. The process should be mild enough that at least 1500 ppm of squalene are retained in the olive oil.

5 The obtained olive oil can be used for the manufacture of spread compositions in which olive oil but not its odour is a desired ingredient. The invention therefore consists in an oil containing spread composition where 5 - 100 wt.% of the oil consists of olive oil and which has no perceivable
10 olive oil odour, characterised in that the composition contains at least 10 ppm of olive oil originating polyphenols.

15

DETAILS OF THE INVENTION

The olive oil containing spreads of the present invention are prepared according to common spread manufacturing process technology. Examples of spread processing can be
20 found e.g. in GB 2281304 and in EP 421 504. Generally spreads are emulsions of a water phase and a fat phase, while the fat phase ususally consists of both liquid oil and a structuring fat. According to the present invention 5 - 100 wt.%, preferably 5 - 95 wt.%, more preferably 10 - 95
25 wt.%, still more preferably 20 - 95 wt.% of the liquid oil of the fat phase of the emulsion consists of olive oil which has been subjected to the mild refining process of the present invention.

30 The invention also comprises a process for the manufacture of any food composition which contains 0.1 - 90 wt.% of olive oil, which process comprises selecting the usual

ingredients, including an olive oil source, and employing the usual processing steps, characterised in that 10 - 100 wt% of the selected olive oil has been obtained according to the process of the invention. The obtained composition
5 may exhibit an olive oil flavour when besides the mild refined olive oil obtained according to the present invention also some ingredient having an olive oil flavour is added as well. So the benefits of a relatively cheap source of refined olive oil and high levels of polyphenol
10 and flavour are combined.

The olive oil flavour source might be e.g. a precious virgin olive oil or a proper substitute, e.g. those obtained by the methods described in WO 97/03566 and EP 852
116.

15

In order to retain after refining a sufficient amount of polyphenols a level of 100 ppm of polyphenols in the starting olive oil is preferred. The starting olive oil preferably is a cheap Lampante oil which usually contains
20 substantial amounts of free fatty acids. Generally, Lampante oil contains from 20 ppm up to 250 ppm of polyphenols.

The refining process not only removes the olive oil odour
25 but also the excess of free fatty acids. High amounts of free fatty acids, even 7 wt.% or more, are often found in Lampante grade olive oils. The mild refining process reduces the fatty acids content to less than 2 wt.%, even down to 0.3 wt.%, which is quite acceptable.

30 Under the mild conditions of the present process the free fatty acids content can be easily reduced to less than 2 wt.%, but obviously the usual low level of 0.1 wt.% is not

attained, unless the starting olive oil is already low in free fatty acids.

Polyphenols are compounds which share a phenolic hydroxyl group. Olive oil originating polyphenols are known to comprise oleuropein, aglycon, tyrosol, hydroxytyrosol and caffeic acid. Their source is olive oil and the olive fruits from which the olive oil is pressed.

The total content of polyphenols in olive oils can be established by standard methods e.g. by the widely accepted colorimetric Gutfinger method as described in J.Am.Oil.Chem.Soc. 1981, 11, pp. 966-968, which method is based on the reaction of a methanolic extract of olive oil and the Folin-Ciocalteu reagent. This method has been used for establishing the polyphenol values mentioned in the present patent specification.

The olive oil is sparged with an inert gas at a temperature up to 250°C under such conditions of sparging gas flow, pressure and process time that the refined oil is free from olive oil odour and still contains at least 1500 ppm of squalene.

Suitable refining conditions are chosen from ranges of relatively moderate sparging temperatures and pressures, particularly 120 - 250°C and 2 - 5 mbar, respectively, using preferably 1 - 15 m³ steam per kg of oil. At 250°C one hour sparging has appeared to be a suitable process time, but time is prolonged to 5 hours when refining takes place at 120°C. Suitable conditions can be established easily by some trials.

Squalene is a high molecular weight ingredient, which occurs at a level of 1500 - 7000 ppm in crude olive oil, including unrefined Lampante oil. Olive oil when refined at traditionally high temperatures contains only little
5 squalene, less than 1500 ppm. An olive oil which has been obtained according to the mild refining process of the present invention has retained squalene at a level of at least 1500 ppm. Experiments have shown that such concentrations can go along with a proper deodorisation.
10 Therefore the content of native squalene has appeared to be a suitable marker for the refining depth of olive oil. A low squalene content points to a great depth of refining carried out under relatively drastic conditions. Such conditions are avoided with the process according to the
15 present invention.

The polyphenols of olive oil are more volatile than squalene, but with the present conditions are retained for a major part in the refined oil.
20 Besides squalene also the above-mentioned unsaponifiables (at least 4500 ppm unsaponifiables) are retained in the mild refined olive oil.
The polyphenol containing spreads prepared according to the invention can be distinguished from known odour-free olive
25 oil containing spreads in that they contain squalene in a concentration which is proportional to the concentration of squalene in the used mild refined olive oil. The invention comprises spreads which contain at least 5 wt.% of olive oil, but which are free from olive oil odour and contain at
30 least 75 ppm of squalene.

Olive oils obtained according to the invention may contain e.g. 90 ppm of polyphenols when refined at 210°C, while the amount drops to 30 ppm when refined at 245°C. Process time and pressure being the same in both cases: about 1.2 hours
5 and about 2.5 mbar. Generally spoken, when a free fatty acid content up to 1 wt.% is allowed, at least 60 wt.% of the polyphenols is retained after refining of the olive oil.

Hence, the olive oil resulting from the found process
10 contains at least 1500 ppm of squalene and is characterized further by the absence of olive oil odour on the one side and by a relatively large amount of native polyphenols on the other side. In practice, absence of olive oil odour is decided when a sensory panel perceives no flavour
15 difference between a composition prepared with an olive oil which has been mild refined according to the invention and a composition prepared with the same ingredients, employing the same processing, but containing an olive oil which has been fully refined in the traditional way.

20 Spreads obtained according to the invention can be recognized because they are characterized by a specific partition of the various types of olive oil polyphenols, which partition is caused by the mild refining conditions which have a different effect on the individual
25 polyphenols. The partition therefore is a kind of fingerprint which reflects the applied refining conditions.

For proper stabilisation of the spreads measures have to be taken in order to prevent flavour reversion. Flavour
30 reversion is known to occur easily when no proper attention

is paid to the removal of metal ions from the olive oil, particularly of iron and copper ions which are notorious catalysts for oil degradation.

According to a preferred embodiment the refining treatment
5 therefore includes measures which aim to reduce the concentration of said metal ions to less than 0.1 ppm. Well known procedures include e.g. the treatment of the olive oil with phosphoric acid and/or with a suitable absorbing bleaching earth, particularly under so-called wet bleaching
10 conditions where the oil has to contain a small amount (0.2 wt.%) of water.

As said before the present mild refining treatment removes the free fatty acids only partially. Under traditional
15 olive oil refining standards the remaining content, amounting to 0.3 - 2 wt.%, is not considered good enough. Fatty acids in such small amounts do not affect, however, product safety or oil taste, while product stability is hardly affected, provided metal ions have been removed.

20

It has appeared that the taste and stability of spreads prepared with the mild refined olive oil complies with the standards set for spreads prepared with traditionally refined olive oil.

25

The invention is illustrated by the following examples:

30

Example 1

OLIVE OIL REFINING

5 Crude olive oil (30 ton), which contained 3.2 wt.% of free fatty acids, 110 ppm of polyphenols and 3000 ppm of squalene was heated up to 90°C and washed by thoroughly mixing with 7 wt.% of soft water of 90°C. The water phase was separated from the olive oil in a self-cleaning
10 centrifuge. By vacuum drying at 50 mbar the olive oil became dry.

The washed and dried olive oil maintained at a temperature of 90°C was first vigorously stirred after addition of an
15 aqueous 50 wt.% phosphoric acid solution (0.8 l per ton of oil). After separation of the aqueous phase by centrifuging, demineralized water was added (0.6 l per ton of oil), followed by the addition of bleaching earth (VOLCANSIL DE22™, 4.5 kg per ton of oil).

20 After 20 min of stirring the pressure was lowered to 40 mbar and stirring was continued for another 20 min. Then the bleaching earth was removed by filtration.

The bleached olive oil was degassed at 100°C and 30 mbar
25 and then gradually heated up to 238°C over one hour. After subsequent steam sparging at 2.3 mbar, spending 9 kg of steam per ton oil, the oil was cooled down to 25°C over one hour and finally cleared by filtration.

Analysis showed that the refined olive oil contained 0.3
30 wt.% of free fatty acids, 65 ppm of polyphenols and 3000 ppm of squalene. A sensory panel assessed the olive oil

flavour. Using a scale with taste scores ranging from 4 (rejected) to 8 (excellent) the panel assigned to the oil a rating of 7, which is almost excellent, while olive oil flavour was hardly perceivable.

5 Even after 4 months of storage no taste defects were noticed.

An average olive oil after been deodorized for one hour at the normal temperature of 255°C contains: 0.1 wt.% of free fatty acids, 1400 ppm of squalene and less than 10 ppm of
10 polyphenols.

Examples 2 - 5

15

OLIVE OIL REFINING

Example 1 was repeated, except that the sparging temperatures were varied as indicated in Table I while exposure times were about the same, but slightly different
20 which explains the variations for polyphenol content.

TABLE I

Ex.	Sparging temperature °C	Polyphenols content ppm	Free fatty acid content wt. %
2	210	88	1.34
3	220	63	0.89
4	230	81	0.52
5	234	62	0.41

25

Example 6

SPREAD CONTAINING MILD REFINED OLIVE OIL

5

A spread was prepared using a water phase and a fat phase having the composition as shown in Table II. The fat phase and the water phase were emulsified by vigorous stirring 10 and pasteurized at 60°C. With a flow of 5-6 kg/h the pasteurized emulsion was conducted through

TABLE II

15

	wt. %
FAT PHASE	60
fat blend ¹	59.55
Hymono 8903 TM	0.08
Bolec ZTD TM	0.08
beta-carotene ²	0.2
flavour	0.09
WATER PHASE	40
skimmed milk powder	1.2
salt	0.09
K-sorbate	0.1
lactic acid	until pH = 5
tap water	balance up to 40

¹ 17% of an interesterified mixture of 65 wt. parts palm oil stearin and 35 wt. parts palm kernel oil and

83% of refined olive oil (containing 1.34 wt.% of free fatty acids and 88 ppm of polyphenols).

² in the form of a 0.4 wt.% beta-carotene solution in sunflower seed oil

5

a Lab VotatorTM line A-A-C. Processing conditions for A1: 800 rpm, exit 20-22°C, A2: 800 rpm, exit 6°C and C (50 ml): 250 rpm.

10 The obtained spread was judged on aspects including flavour, colour, taste, taste stability, spreadability, consistence and stability of consistence. The spread exhibited excellent quality in all aspects.

A comparison spread was prepared under identical conditions
15 and with the same ingredients, except that an olive oil was chosen, qualified as pure olive oil, which was obtained by traditional refining at a relatively high temperature. However, a sensory panel could perceive a taste difference between both spreads, neither immediately after
20 preparation, nor after 5 weeks of storage.

CLAIMS

1. An oil containing spread composition, where 5 - 100 wt.% of the oil consists of olive oil and which has no perceivable olive oil odour, characterised in that the composition contains at least 10 ppm, preferably at least 50 ppm, more preferably at least 200 ppm of olive oil originating polyphenols.
2. A spread according to claim 1 containing at least 5 wt.% of olive oil and at least 75 ppm of squalene.
3. A process for refining an olive oil which process is characterised in that the oil is sparged with an inert gas at a temperature up to 250°C under such conditions of sparging gas flow, pressure and process time that the refined oil is free from olive oil odour and contains at least 1500 ppm of squalene.
4. A process according to claim 3, characterized in that the olive oil to be refined, the source oil, contains at least 100 ppm of polyphenols.
5. A process according to claim 3, characterized in that the refining conditions are such that at least 60 wt.% of the polyphenols present in the source oil are retained.
6. A process according to claim 3, characterized in that the sparging temperature is 120 - 250°C.

7. A process according to claim 3, characterized in that the sparging pressure is 2 - 5 mbar.
8. A process for the manufacture of a food composition which contains 0.1 - 90 wt.% of olive oil, comprising selecting the usual ingredients, including an olive oil source, and employing the usual processing steps, characterised in that 10 - 100 wt% of the olive oil has been obtained according to any one of claims 3 - 7.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/07797

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C11B3/14 A23D7/00 A23D9/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C11B A23D A23L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 94 12596 A (UNILEVER PLC ;UNILEVER NV (NL)) 9 June 1994 see page 4, line 10 - line 28 see page 5, line 20 - line 30 see claims 1,7,11-14 ---	1,3,6-8
Y	WO 97 32947 A (COMITER TRADING & SERVICES S R ;BALDI ALESSANDRO (IT); ROMANI ANNA) 12 September 1997 see page 3, line 17 - line 21 see claims 1,2,7,10,20 ---	1
Y	EP 0 304 354 A (SYNTHELABO) 22 February 1989 see page 2, line 24 - line 31 see claims 1,2,6-8 ---	3,6-8
	--- -/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

21 May 1999

Date of mailing of the international search report

04/06/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Dekeirel, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/07797

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, A	EP 0 849 353 A (UNILEVER PLC ; UNILEVER NV (NL)) 24 June 1998 see claim 1 ---	1
A	GB 1 511 953 A (ASAHI DENKA KOGYO KK) 24 May 1978 see page 2, line 122 - page 3, line 8 see claims 1, 4, 6 ---	3, 6, 7
A	EP 0 724 874 A (NESTLE SA) 7 August 1996 see example 1 -----	3, 6, 7

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/07797

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9412596 A	09-06-1994	AT 152761 T AU 672858 B AU 5624694 A DE 69310543 D DE 69310543 T DK 672096 T EP 0672096 A ES 2102806 T ZA 9308754 A	15-05-1997 17-10-1996 22-06-1994 12-06-1997 04-09-1997 03-11-1997 20-09-1995 01-08-1997 23-05-1995
WO 9732947 A	12-09-1997	IT F1960041 A EP 0896612 A	05-09-1997 17-02-1999
EP 0304354 A	22-02-1989	FR 2618449 A AT 68819 T AU 599502 B AU 1972088 A CA 1335111 A DE 3865777 A DK 413388 A FI 883485 A,B, JP 1048897 A NO 175009 B PT 88084 A,B	27-01-1989 15-11-1991 19-07-1990 27-01-1989 04-04-1995 28-11-1991 24-01-1989 24-01-1989 23-02-1989 09-05-1994 30-06-1989
EP 0849353 A	24-06-1998	NONE	
GB 1511953 A	24-05-1978	JP 51109906 A	29-09-1976
EP 0724874 A	07-08-1996	AU 692031 B AU 4330096 A CA 2168495 A JP 8245367 A NO 960430 A NZ 280918 A US 5653966 A	28-05-1998 08-08-1996 03-08-1996 24-09-1996 05-08-1996 28-10-1996 05-08-1997